

Patient Handout

Patient Education
lurasidone oral

LURASIDONE - ORAL

(loo-RAS-i-done)

COMMON BRAND NAME(S): Latuda

WARNING: There may be a slightly increased risk of serious, possibly fatal side effects (such as stroke, heart failure, fast/irregular heartbeat, pneumonia) when this medication is used by older adults with dementia. This medication is not approved for the treatment of dementia-related behavior problems. Discuss the risks and benefits of this medication, as well as other effective and possibly safer treatments for dementia-related behavior problems, with the doctor. Lurasidone is used to treat certain mental/mood disorders (such as schizophrenia, depression associated with bipolar disorder). Drugs used to treat depression can help prevent suicidal thoughts/attempts and provide other important benefits. However, studies have shown that a small number of people (especially people younger than 25) who take drugs to treat depression may experience worsening depression, other mental/mood symptoms, or suicidal thoughts/attempts. It is very important to talk with the doctor about the risks and benefits of drugs used to treat depression (especially for people younger than 25), even if treatment is not for a mental/mood condition. Tell the doctor right away if you notice worsening depression/other psychiatric conditions, unusual behavior changes (including possible suicidal thoughts/attempts), or other mental/mood changes (including new/worsening anxiety, panic attacks, trouble sleeping, irritability, hostile/angry feelings, impulsive actions, severe restlessness, very rapid speech). Be especially watchful for these symptoms when a new drug to treat depression is started or when the dose is changed.

USES: This medication is used to treat certain mental/mood disorders (such as schizophrenia, depression associated with bipolar disorder). Lurasidone belongs to a class of drugs known as

atypical antipsychotics. It works by helping to restore the balance of certain natural substances in the brain.

HOW TO USE: Read the Medication Guide provided by your pharmacist before you start taking lurasidone and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth with food as directed by your doctor, usually once daily. The dosage is based on your medical condition, response to treatment, age, and other medications you may be taking. Be sure to tell your doctor and pharmacist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Take this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Keep taking this medication exactly as prescribed, even if you are feeling better and thinking more clearly. Do not increase your dose or take this drug more often than prescribed. Your symptoms will not improve any faster, and your risk of side effects will increase. Do not stop taking this medication without consulting your doctor. Avoid eating grapefruit or drinking grapefruit juice while using this medication unless your doctor or pharmacist says you may do so safely. Grapefruit can increase the chance of side effects with this medicine. Ask your doctor or pharmacist for more details. Tell your doctor if your condition does not improve or if it worsens. It may take several weeks before you feel the full benefit of this medication.

SIDE EFFECTS: See also Warning section. Drowsiness, dizziness, lightheadedness, nausea, shaking, weight gain, mask-like facial expression, inability to keep still, and agitation may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Your doctor may order another medication to lessen these effects. This medication may cause a serious drop in blood pressure, especially when starting this medication. Dizziness and lightheadedness can increase the risk of falling. Get up slowly when rising from a sitting or lying position. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: drooling/trouble swallowing, fainting, interrupted breathing during sleep. Rarely, this medication may cause face/muscle twitching and uncontrollable movements (tardive dyskinesia). In some cases, this condition may be permanent. Tell your doctor right away if you develop any uncontrollable movements such as lip smacking, mouth puckering, tongue thrusting, chewing, or unusual arm/leg movements. This drug may rarely make your blood sugar rise, which can cause or worsen diabetes. Weight gain from this drug may increase the risk of this side effect. Tell your doctor right away if you have symptoms of high blood sugar such as increased thirst/urination. If you already have diabetes, check your blood sugar regularly as directed and share the results with your doctor. Your doctor may need to adjust your diabetes medication, exercise program, or diet. This medication may also cause a rise in your blood fat levels (cholesterol and triglycerides). These increases, along with diabetes and weight gain, may increase your risk for developing heart disease. Discuss the risks and benefits of treatment with your doctor. In rare cases, lurasidone may increase your level of a certain substance made by the body (prolactin). For females, this increase in prolactin may result in unwanted breast milk, missed/stopped periods, or difficulty becoming pregnant. For males, it may result in decreased sexual ability, inability to produce sperm, or enlarged breasts. If you develop any of these symptoms, tell your doctor right away. Rarely, males may have a painful or prolonged erection lasting 4 or more hours. If this occurs, stop using this drug and get medical help right away, or permanent problems could occur. Get medical help right away if you have any very serious side effects, including: seizure. This medication may rarely cause a very serious condition called neuroleptic malignant syndrome (NMS). Get medical help right away if you have any of the following symptoms: fever, muscle stiffness/pain/tenderness/weakness, severe tiredness, severe confusion, sweating, fast/irregular heartbeat, dark urine, signs of kidney problems (such as change in the amount of urine). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: fever, swollen lymph nodes, rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US -Call your doctor for medical advice about

side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

PRECAUTIONS: See also Warning section. Before taking lurasidone, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney problems, liver problems, stroke, breast cancer, diabetes (including family history), obesity, low blood pressure, seizures, low white blood cell count, dementia (such as Alzheimer's Disease), trouble swallowing, breathing trouble during sleep (sleep apnea). This drug may make you dizzy or drowsy. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Avoid alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This medication may make you sweat less, making you more likely to get heat stroke. Avoid doing things that may cause you to overheat, such as hard work or exercise in hot weather, or using hot tubs. When the weather is hot, drink a lot of fluids and dress lightly. If you overheat, quickly look for a place to cool down and rest. Get medical help right away if you have a fever that does not go away, mental/mood changes, headache, or dizziness. Older adults may be more sensitive to the side effects of this drug, especially drowsiness, dizziness, and lightheadedness. Drowsiness, dizziness, and lightheadedness can increase the risk of falling. During pregnancy, this medication should be used only when clearly needed. Babies born to mothers who have used this drug during the last 3 months of pregnancy may rarely develop symptoms including muscle stiffness or shakiness, drowsiness, feeding/breathing difficulties, or constant crying. If you notice any of these symptoms in your newborn especially during their first month, tell the doctor right away. Since untreated mental/mood problems (such as schizophrenia, bipolar disorder, depression) can be a serious condition, do not stop taking this medication unless directed by your doctor. If you are planning pregnancy, become pregnant, or think you may be pregnant, immediately discuss with your doctor the benefits and risks of using this medication during pregnancy. It is unknown if this medication passes into breast milk. Consult your doctor before breastfeeding.

DRUG INTERACTIONS: See also How To Use section. Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Other medications can affect the removal of lurasidone from your body, which may affect how lurasidone works. Examples include cobicistat, diltiazem, ritonavir, telithromycin, azole antifungals (such as ketoconazole, itraconazole), certain HIV protease inhibitors (such as nelfinavir), macrolide antibiotics (such as clarithromycin), rifamycins (such as rifampin, rifabutin), antidepressants (such as fluoxetine, paroxetine, nefazodone), among others. Tell your doctor or pharmacist if you are taking other products that cause drowsiness such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), muscle relaxants (such as carisoprodol, cyclobenzaprine), or antihistamines (such as cetirizine, diphenhydramine). Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause dizziness or drowsiness. Ask your pharmacist about using those products safely.

OVERDOSE: If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call 1-800-222-1222. Canada residents can call 1-844-764-7669.

NOTES: Do not share this medication with others. Lab and/or medical tests (such as blood sugar, complete blood count, blood pressure, weight, cholesterol/triglyceride levels) may be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details.

MISSED DOSE: If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up.

STORAGE: Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Information last revised May 2024. Copyright(c) 2024 First Databank, Inc.

IMPORTANT: HOW TO USE THIS INFORMATION: This is a summary and does NOT have all possible information about this product. This information does not assure that this product is safe, effective, or appropriate for you. This information is not individual medical advice and does not substitute for the advice of your health care professional. Always ask your health care professional for complete information about this product and your specific health needs.

TOP PICKS FOR YOU

Get the Free Medscape App

Download on the App Store Get it on Google Play

Or, send yourself a link to download the app later.

Email

Please use a valid email address

Don't show me this again

Your email has been sent

Resend Email

Unlimited Access & Benefits in Your Pocket:

Offline access to the Medscape drug database

Easy navigation of drug information

Interaction Checker, Pill Identifier, and more clinical tools